



## The HIV response beyond 2030: preparing for decades of sustained HIV epidemic control in eastern and southern Africa



Published Online  
May 20, 2024  
[https://doi.org/10.1016/S0140-6736\(24\)00980-2](https://doi.org/10.1016/S0140-6736(24)00980-2)

The UN global goal to “end AIDS as a public health threat by 2030”<sup>1,2</sup> has motivated remarkable progress in eastern and southern African countries most affected by HIV. In these countries since 2010, new HIV infections have decreased by an estimated 57% and AIDS-related deaths by 58%.<sup>3</sup> Targeted 90% reductions between 2010 and 2030 are in reach for some countries in eastern and southern Africa that are also on track to attain the UNAIDS 95-95-95 HIV care cascade targets.<sup>3-6</sup>

The 2015 UNAIDS–*Lancet* Commission envisioned “ending AIDS” as achieving disease control, recognising that long-term intervention measures would be needed to maintain the lowered rates of new HIV infections and AIDS-related deaths.<sup>7,8</sup> To sustain HIV epidemic control beyond 2030, countries with high burdens of HIV will need to continue steadily further reducing new HIV infections over coming decades, eventually to below a globally applicable threshold, such as below one new infection per 10 000 HIV-negative population.<sup>9,10</sup> Continued declines in HIV infections are crucial to contain long-term resources required for providing HIV treatment and to avoid risk of resurgent HIV transmission. For countries reaching the UNAIDS 95-95-95 targets, mathematical model projections suggest a further 20% reduction in new HIV infections every 5 years is an ambitious but attainable target to guide prevention strategies.<sup>11,12</sup> Where larger care cascade gaps remain, quickly increasing HIV treatment coverage would rapidly reduce population viraemia, enabling steeper HIV incidence declines.<sup>13-15</sup>

Through a meeting series convened by UNAIDS, the Post-2030 HIV Response Working Group reviewed progress in the HIV response, the evolving nature of the epidemic, how to define long-term epidemic control beyond 2030, and the key programmes, policies, and surveillance required to ensure it is sustained.<sup>9,11</sup> Here, we identify four essential priorities to sustain HIV epidemic control in countries in eastern and southern Africa with large HIV epidemics and successful HIV programmes.<sup>9</sup>

First, effective HIV treatment is the cornerstone of success. Even with continued success in reducing new HIV infections, it is estimated that the 21 million people living with HIV today in eastern and southern Africa will decline only gradually to about 13–17 million people

living with HIV by 2050,<sup>9,16</sup> underscoring the need for long-term programmes delivering lifelong antiretroviral therapy. Maintaining extremely high treatment coverage and undetectable viral load is essential for the health of people living with HIV and reducing transmission, representing the powerful alignment of individual and population health outcomes embodied by the U=U (Undetectable=Untransmittable) public health message.<sup>17</sup> Providing antiretroviral therapy will constitute the majority of future resources for HIV programmes. Disruption to supply chains or delivery could precipitate immediate rapid rises in AIDS-related deaths and new HIV infections,<sup>18</sup> while deterioration in HIV treatment continuation or effectiveness at durable viral suppression, for example through increasing drug resistance, risks slowing declines in HIV incidence and thereby increasing future resource requirements for HIV care and treatment.<sup>19</sup>

Second, it is also important to ensure timely HIV diagnosis. HIV testing programmes should transition focus from the proportion of HIV positive people aware of their status in the 95-95-95 targets to ensuring short time to diagnosis which enables rapid viral suppression. HIV testing is fairly inexpensive and should be easily accessible to anyone, increasingly through self-testing,<sup>20</sup> with frequent testing encouraged among people with increased risk of exposure to HIV acquisition.

Third, HIV prevention approaches need to adapt with evolving individual needs and preferences to ensure continued use of prevention methods at levels that keep HIV infections low. The diffuse nature of HIV transmission in contemporary African HIV epidemics<sup>21</sup> necessitates strategies that engage large populations with moderate HIV risk in effective, easily accessible, and affordable prevention options, such as use of condoms and voluntary medical male circumcision. People with increased exposure to HIV acquisition, including some young people, need more intensive prevention choices, such as pre-exposure prophylaxis.<sup>22</sup> Deterioration in HIV testing or prevention threatens epidemic control through decelerating or stalling HIV incidence declines,<sup>19</sup> which would be expected to only become apparent 5–10 years later.

Fourth, comprehensive HIV services for key populations need to be maintained for the long term,<sup>23</sup> including access to new antiviral-based prevention technologies. New HIV infections among key populations, including sex workers, gay men and other men who have sex with men, people who inject drugs, transgender people, and people in prisons, are about 9% of all new HIV infections in eastern and southern Africa, but occur at rates four to ten times higher than in all adults.<sup>24</sup> Theoretical epidemic dynamics suggest the proportion of HIV infections among key populations could increase as overall infections decline,<sup>25</sup> but this rise is not inevitable.<sup>26</sup> Services that meet the distinct prevention needs of key populations address health equity and human rights for key populations and ensure long-term epidemic control.

Shifting focus from rapid intervention scale-up to implementing resilient decades-long HIV programmes entails myriad new challenges for the HIV response. Vertical HIV management and delivery systems, while effective, are fragile to changing priorities. Such systems must maintain effectiveness while being integrated into local, national, and regional structures<sup>27</sup> to ensure durability of HIV services. Societally, the impacts of HIV programmes have been enabled by successfully addressing societal and structural barriers to accessing HIV services.<sup>28</sup> Current legislative attacks on human rights that criminalise populations at risk of HIV,<sup>29,30</sup> such as gay men and other men who have sex with men,<sup>31</sup> threaten the ability to ensure supportive legal environments, gender equality, ending stigma and discrimination, and the multisectoral coordination required for sustained future progress in controlling the HIV epidemic in countries in eastern and southern Africa.

HIV treatment programmes need to adapt to the changing health needs of ageing populations with HIV. In eastern and southern African countries, the median age of people living with HIV will increase from 32 years in 2010 to 59 years by 2050.<sup>9</sup> Maintaining effective HIV treatment requires a continued focus on improving care quality,<sup>32</sup> integration with primary care,<sup>33,34</sup> and addressing access barriers imposed by stigma and discrimination.<sup>35</sup> Equally, as HIV epidemics recede, high awareness and motivation for HIV prevention and testing could become more challenging, especially among young people unfamiliar with the height of the HIV/AIDS emergency. Policies and programmes will need more focused attention to ensuring HIV prevention, testing, and treatment equitably

reach mobile, marginalised, and socioeconomically disadvantaged populations. As evidenced throughout the HIV/AIDS response,<sup>36</sup> empowered communities can safeguard success through delivering person-centred HIV services, guiding priorities and improvement in the quality of services, and holding governments accountable during HIV service integration and management transitions.

Lastly, ongoing innovation will provide new tools that can support sustaining epidemic control. Ensuring scalable and affordable access to medicines and commodities for the countries and communities most affected by HIV will maximise the impact of new long-acting antiretrovirals and pre-exposure and post-exposure prophylaxis. Digital health information systems and new point-of-care diagnostics for HIV viral load and resistance monitoring,<sup>37</sup> if resourced and implemented at scale, will unlock more convenient person-centred HIV service delivery models, decongest health facilities, and facilitate efficient surveillance to identify and mitigate emerging threats to epidemic control.

Considerable gains have been made in responding to HIV in eastern and southern Africa, but current progress is fragile. As 2030 approaches, the focus of—and terminology used in—regional and national HIV agendas should evolve from referring to the “end of AIDS” to building momentum for decades of sustained and resilient HIV epidemic control. These efforts need to include concerted actions towards integrated long-term services for millions of people living with HIV, minimising new HIV infections, and confronting HIV stigma and discrimination.

JWI-E's institution receives funding from the Bill & Melinda Gates Foundation (BMGF), the Joint United Nations Programme on HIV/AIDS (UNAIDS), and the US National Institutes of Health (NIH) and has received consulting fees from BAO Systems and support to attend meetings from UNAIDS, BMGF, the International AIDS Society, and the South African Centre for Epidemiological Modelling and Analysis (SACEMA). CF's institution receives funding from BMGF for the PANGEA2-HIV consortium and received support from UNAIDS to attend meetings in Boston and Botswana related to this work. HA received support from UNAIDS and BMGF to attend a meeting in Botswana related to this work. JSt's institution receives funding from UNAIDS and received support from UNAIDS and BMGF to attend a meeting in Botswana related to this work. PG-F received support from Imperial College London to support this work and to attend the meeting in Botswana and has received consultancy contracts from UNAIDS for other projects. LA-D, OE, and SMi received support from UNAIDS to attend meetings in Boston and Botswana related to this work. RA, SMa, SLB, and WLK received support from UNAIDS to attend a meeting in Botswana related to this work. RG's institution receives funding from UNAIDS and BMGF. MKG's institution receives funding from BMGF. AA, MRM, and GPG are employees of BMGF. AB's institution receives funding from NIH, BMGF, the New York City Department of Health and Mental Hygiene, and the Foundation for Innovative New Diagnostics (FIND) and has received consulting fees from Gates Ventures. HM received support from BMGF for work presented at meetings related to this Comment. JRH receives funding from BMGF and was

partly supported by the MeSH Consortium, based at the London School of Hygiene & Tropical Medicine with a grant funded by BMGF (INV-007055). GM-R's institution receives funding from BMGF, FIND, and the United States Agency for International Development (USAID). WJMP received consulting fees from WHO. ANP's institution receives funding from BMGF, the Wellcome Trust, the European Union (EU), NIH, and the National Institute for Health and Care Research (NIHR). VC's institution receives funding from BMGF, NIHR, USAID, Unitaid, UK Research and Innovation (UKRI), and the Medical Research Council (MRC) and has received consulting fees from WHO. DK has received consulting fees from University College London and Massachusetts General Hospital. MREHP has received consulting fees from the HIV Modelling Consortium. LB-M's institution receives funding from BMGF. SPR is an employee of Airfinity Ltd. LKW's institution receives funding from Wellcome Trust, NIHR, has received consulting fees from WHO and Pacific Life Re and has received honoraria from *The Lancet Infectious Diseases* and Luxembourg National Research Fund and has participated on a Research Steering Group for the Wellcome Trust. All the other authors declare no competing interests.

Copyright © 2024 World Health Organization. Published by Elsevier Ltd. All rights reserved

### The Post-2030 HIV Response Working Group jeaton@hsph.harvard.edu

The Post-2030 HIV Response Working Group, c/o Data for Impact Department, UNAIDS, Geneva, Switzerland

The members of the The Post-2030 HIV Response Working Group are Adam Akullian, Ruth Akulu, Gambo Aliyu, Florence Anam, Anne-Claire Guichard, Helen Ayles, Rachel Baggaley, Lovleen Banshi-Matharu, Solange L Baptiste, Anna Bershteyn, Valentina Cambiano, Austin Carter, Nafisah Chotun, Daniel T Citron, Siobhan Crowley, Shona Dalal, Olanrewaju Edun, Christophe Fraser, Alison P Galvani, Geoffrey P Garnett, Robert Glaubius, Peter Godfrey-Faussett, M Kate Grabowski, Glenda E Gray, James R Hargreaves, Jeffrey W Imai-Eaton, Leigh F Johnson, David Kaftan, Joseph Kagaayi, Edward Kataika, Nduku Kilonzo, Wilford L Kirungi, Eline L Korenromp, Mach-Houd Kouton, Lucie Abeler-Dörner, Mary Mahy, Tara D Mangal, Rowan Martin-Hughes, Samuel Matsikure, Gesine Meyer-Rath, Sharmistha Mishra, Mpho Mmehesi, Abdulaziz Mohammed, Haroon Moolla, Michelle R Morrison, Sikhulile Moyo, Edinah Mudimu, Mbulawa Mugabe, Maurine Murenga, Joyce Ng'ang'a, Yewande Olaifa, Andrew N Phillips, Michael R E H Pickles, William J M Probert, Dinah Ramaabya, Stefan P Rautenbach, Paul Revill, Ani Shakarishvili, Robert Sheneberger, Jennifer Smith, Christine Stegling, John Stover, Frank Tanser, Isaac Taramusi, Debra ten Brink, Lilith K Whittles, and Irum Zaidi. The affiliations of the group members are listed in the appendix.

- 1 UNAIDS. Fast-track: ending the AIDS epidemic by 2030. 2014. [http://www.unaids.org/sites/default/files/media\\_asset/C2686\\_WAD2014report\\_en.pdf](http://www.unaids.org/sites/default/files/media_asset/C2686_WAD2014report_en.pdf) (accessed May 1, 2024).
- 2 African Union. Catalytic framework to end AIDS, tuberculosis and eliminate malaria in Africa by 2030. 2016. [https://au.int/sites/default/files/pages/32904-file-catalytic\\_framework\\_8pp\\_en\\_hires.pdf](https://au.int/sites/default/files/pages/32904-file-catalytic_framework_8pp_en_hires.pdf) (accessed May 1, 2024).
- 3 UNAIDS. The path that ends AIDS: UNAIDS Global AIDS Update 2023. 2023. [https://thepath.unaids.org/wp-content/themes/unaids2023/assets/files/2023\\_report.pdf](https://thepath.unaids.org/wp-content/themes/unaids2023/assets/files/2023_report.pdf) (accessed May 1, 2024).
- 4 Zimbabwe Ministry of Health and Child Care. Zimbabwe Population-Based HIV Impact Assessment ZIMPHIA 2020: summary sheet. December, 2020. [https://phia.icap.columbia.edu/wp-content/uploads/2020/11/ZIMPHIA-2020-Summary-Sheet\\_Web.pdf](https://phia.icap.columbia.edu/wp-content/uploads/2020/11/ZIMPHIA-2020-Summary-Sheet_Web.pdf) (accessed May 1, 2024).
- 5 Malawi Ministry of Health. Malawi Population-based HIV Impact Assessment MPHIA 2020–2021: summary sheet. March, 2022. [https://phia.icap.columbia.edu/wp-content/uploads/2022/03/110322\\_MPHIA\\_Summary-sheet-English.pdf](https://phia.icap.columbia.edu/wp-content/uploads/2022/03/110322_MPHIA_Summary-sheet-English.pdf) (accessed May 1, 2024).
- 6 Eswatini Ministry of Health and Central Statistical Office. Eswatini Population-Based HIV Impact Assessment SHIMS3 2021: summary sheet. December, 2022. [https://phia.icap.columbia.edu/wp-content/uploads/2022/12/53059\\_14\\_SHIMS3\\_Summary-sheet-Web.pdf](https://phia.icap.columbia.edu/wp-content/uploads/2022/12/53059_14_SHIMS3_Summary-sheet-Web.pdf) (accessed May 1, 2024).
- 7 Dowdle WR. The principles of disease elimination and eradication. *Bull World Health Organ* 1998; **76**: 22.
- 8 Piot P, Abdool Karim SS, Hecht R, et al. Defeating AIDS—advancing global health. *Lancet* 2015; **386**: 171–218.
- 9 UNAIDS. Charting the post-2030 AIDS pandemic and response: what is needed to sustain gains addressing HIV in eastern and southern Africa for the long term. May 20, 2024. <https://jointsiwg.unaids.org/wp-content/uploads/2024/05/2023-Charting-the-post-2030-AIDS-response.pdf> (accessed May 20, 2024).
- 10 Galvani AP, Pandey A, Fitzpatrick MC, Medlock J, Gray GE. Defining control of HIV epidemics. *Lancet HIV* 2018; **5**: e667–70.
- 11 UNAIDS. Describing “the End of AIDS as a Public Health Threat”: Final Report of a Technical Working Meeting held at the Harvard T H Chan School of Public Health, Boston, MA, USA. July 20–21, 2023. [https://jointsiwg.unaids.org/wp-content/uploads/2024/02/Describing\\_End\\_of\\_AIDS\\_Harvard\\_Technical\\_Mtg\\_Edited\\_Layout\\_26\\_Feb\\_2024.pdf](https://jointsiwg.unaids.org/wp-content/uploads/2024/02/Describing_End_of_AIDS_Harvard_Technical_Mtg_Edited_Layout_26_Feb_2024.pdf) (accessed May 13, 2024).
- 12 Moolla H, Phillips A, ten Brink D, et al. A quantitative assessment of the consistency of projections from five mathematical models of the HIV epidemic in South Africa: a model comparison study. *BMC Public Health* 2023; **23**: 2119.
- 13 Hladik W, Stupp P, McCracken SD, et al. The epidemiology of HIV population viral load in twelve sub-Saharan African countries. *PLoS One* 2023; **18**: e0275560.
- 14 Larmarange J, Bachanas P, Skalland T, et al. Population-level viremia predicts HIV incidence at the community level across the Universal Testing and Treatment Trials in eastern and southern Africa. *PLOS Glob Public Health* 2023; **3**: e0002157.
- 15 Tanser F, Vandormael A, Cuadros D, et al. Effect of population viral load on prospective HIV incidence in a hyperendemic rural African community. *Sci Transl Med* 2017; **9**: eaam8012.
- 16 Stover J, Glaubius R, Teng Y, et al. Modeling the epidemiological impact of the UNAIDS 2025 targets to end AIDS as a public health threat by 2030. *PLoS Med* 2021; **18**: e1003831.
- 17 Eisinger RW, Dieffenbach CW, Fauci AS. HIV viral load and transmissibility of HIV infection: Undetectable equals Untransmittable. *JAMA* 2019; **321**: 451–52.
- 18 Jewell BL, Mudimu E, Stover J, et al. Potential effects of disruption to HIV programmes in sub-Saharan Africa caused by COVID-19: results from multiple mathematical models. *Lancet HIV* 2020; **7**: e629–40.
- 19 Rautenbach SP, Whittles LK, Meyer-Rath G, et al. Future HIV epidemic trajectories in South Africa and projected long-term consequences of reductions in general population HIV testing: a mathematical modelling study. *Lancet Public Health* 2024; **9**: e218–30.
- 20 WHO. WHO recommends optimizing HIV testing services. 2024. <https://www.who.int/news/item/22-07-2023-who-recommends-optimizing-hiv-testing-services> (accessed April 25, 2024).
- 21 Otecko N, Hall M, Di Lauro F, et al. HIV transmission clusters in Zambia are smaller than in Europe: evidence from the HPTN-071 (PopART) study, BEEHIVE study, and individual-based models. IAS 2023, the 12th IAS Conference on HIV Science. Brisbane, Australia. July 23, 2023. [https://plus.iasociety.org/sites/default/files/2023-11/e-poster\\_885.pdf](https://plus.iasociety.org/sites/default/files/2023-11/e-poster_885.pdf) (accessed May 1, 2024).
- 22 UNAIDS. End Inequalities. End AIDS. Global AIDS Strategy 2021–2026. 2021. <https://www.unaids.org/en/resources/documents/2021/2021-2026-global-AIDS-strategy> (accessed May 13, 2024).
- 23 WHO. Consolidated guidelines on HIV, viral hepatitis and STI prevention, diagnosis, treatment and care for key populations. Geneva: World Health Organization, 2022.
- 24 Korenromp EL, Sabin K, Stover J, et al. New HIV infections among key populations and their partners in 2010 and 2022, by world region: a multisources estimation. *J Acquir Immune Defic Syndr* 2024; **95**: e34–45.
- 25 Garnett GP. Reductions in HIV incidence are likely to increase the importance of key population programmes for HIV control in sub-Saharan Africa. *J Int AIDS Soc* 2021; **24** (suppl 3): e25727.
- 26 Jones HS, Anderson RL, Cust H, et al. HIV incidence among women engaging in sex work in sub-Saharan Africa: a systematic review and meta-analysis. *medRxiv* 2023; published online Oct 18. <https://doi.org/10.1101/2023.10.17.23297108> (preprint).
- 27 Africa Centres for Disease Control and Prevention. Africa CDC strategic plan 2023–2027. 2023. <https://africacdc.org/download/africa-cdc-strategic-plan-2023-2027/> (accessed May 1, 2024).
- 28 Stangl AL, Pliakas T, Izazola-Licea JA, et al. Removing the societal and legal impediments to the HIV response: an evidence-based framework for 2025 and beyond. *PLoS One* 2022; **17**: e0264249.
- 29 Beyrer C, Kamarulzaman A, Isbell M, et al. Under threat: the International AIDS Society–Lancet Commission on Health and Human Rights. *Lancet* 2024; **403**: 1374–418.

See Online for appendix

- 30 UNDP, UNAIDS. Preventing and responding to an HIV-related human rights crisis. 2024. [https://www.unaids.org/sites/default/files/media\\_asset/hiv-related-human-rights-crisis\\_en.pdf](https://www.unaids.org/sites/default/files/media_asset/hiv-related-human-rights-crisis_en.pdf) (accessed April 25, 2024).
- 31 Dyer O. Surge in anti-gay laws in Africa imperil progress on HIV/AIDS, experts say. *BMJ* 2024; **384**: q599.
- 32 Safreed-Harmon K, Anderson J, Azzopardi-Muscat N, et al. Reorienting health systems to care for people with HIV beyond viral suppression. *Lancet HIV* 2019; **6**: e869–77.
- 33 Goldstein D, Salvatore M, Ferris R, Phelps BR, Minior T. Integrating global HIV services with primary health care: a key step in sustainable HIV epidemic control. *Lancet Glob Health* 2023; **11**: e1120–24.
- 34 WHO. Primary health care and HIV: convergent actions. Policy considerations for decision-makers. Geneva: World Health Organization, 2023.
- 35 UNAIDS. The path that ends AIDS: UNAIDS Global AIDS Update 2023. 2023. <https://www.unaids.org/en/resources/documents/2023/global-aids-update-2023> (accessed May 1, 2024).
- 36 UNAIDS. Let communities lead: World AIDS Day report 2023. 2023. [https://www.unaids.org/sites/default/files/media\\_asset/2023WADreport\\_en.pdf](https://www.unaids.org/sites/default/files/media_asset/2023WADreport_en.pdf) (accessed May 1, 2024).
- 37 Bonsall D, Golubchik T, de Cesare M, et al. A comprehensive genomics solution for HIV surveillance and clinical monitoring in low-income settings. *J Clin Microbiol* 2020; **58**: e00382–20.